Asymmetries and Tau Polarization at LEP

Günter Duckeck
LMU Munich
WIN 99, Cape Town, January 1999

I. Introduction

II. Lepton forward–backward asymmetries

III. Quark forward–backward asymmetries

IV. Tau polarization

V. Summary

Introduction

Asymmetry at LEP caused by small parity violation in leptonic Z^0 couplings:



$\rightarrow$ Z^0 polarized: $\mathcal{A}_e = \frac{2\,\hat{g}_v^e\,\hat{g}_a^e}{(\hat{g}_v^e)^2+(\hat{g}_a^e)^2}$.

Similar in decay:



$\rightarrow$ final state polarized $\mathcal{A}_f = \frac{2\,\hat{g}_v^f\,\hat{g}_a^f}{(\hat{g}_v^f)^2+(\hat{g}_a^f)^2}$.

Observables

- Residual polarization of final state (measurable in τ decays)

$$\langle P_\tau \rangle = \frac{\sigma_r - \sigma_l}{\sigma_r + \sigma_l} \approx -\mathcal{A}_\tau$$

- Polarization asymmetry

$$A_{\rm FB}^{\rm pol} = \frac{(\sigma_r^F - \sigma_l^F) - (\sigma_r^B - \sigma_l^B)}{\sigma_{tot}} \approx -\frac{3}{4}\mathcal{A}_e$$

- Forward-backward charge asymmetry

$$A_{\rm FB}^f = \frac{\sigma_f^F - \sigma_f^B}{\sigma_f^F + \sigma_f^B} = \frac{3}{4}\mathcal{A}_e \cdot \mathcal{A}_f$$

Observables related to $\sin^2\theta^{\,l}_{\rm eff}$:

$$\mathcal{A}_f = \frac{2(1-4|Q_f|\sin^2\theta^{\,l}_{\rm eff})}{1+(1-4|Q_f|\sin^2\theta^{\,l}_{\rm eff})^2}$$



ISR, γ, γZ and $\sqrt{s}$ modify pure Z^0 effects:



Lepton charge asymmetries

Conceptually 'cleanest' measurement at LEP (in particular $\mu^+\mu^-$)

- clear separation of lepton species



- unambiguous charge identification

- A_{FB}^{ℓ} is ratio of numbers
  $\rightarrow$ luminosity uncertainty and most slection systematics cancel
- t-channel diagrams complicate e^+e^- channel





A_{FB} determined in maximum likelihood fit to differential distribution

$$\frac{1}{\sigma} = \frac{d\sigma}{d\cos\theta} = \frac{3}{8}\left(1+\cos^2\theta\right) + A_{\mathrm{FB}}\cos\theta$$

Systematics

Systematics of A_{FB} measurement

- charge & efficiency symmetry of detector & selection
- fiducial cut on angular range (e^+e^- t-channel)
- background (e^+e^- for $\tau^+\tau^-$)
- physics (asymmetric acceptance for e^+e^-, τ polarization)

in general small, well below statistics.

$A_{\mathrm{FB}}^{0,\ell}$ determined in global lineshape fit to σ and asymmetries, accounts for all effects of ISR, γ, γZ and $\sqrt{s}$.

t-channel correction done with ALIBABA/TOPAZ.
Introduces additional systematics of ≈ 0.0013 and correlations with $\sigma_{e^+e^-}$.
$\rightarrow$ Common to all experiments

Results


OPAL Preliminary
ee
1990
1991
1992
1993
1994
1995
Asymmetry
Data - Fit
√s (GeV)


OPAL Preliminary
μμ
1990
1991
1992
1993
1994
1995
Asymmetry
Data - Fit
√s (GeV)


OPAL Preliminary
ττ
1990
1991
1992
1993
1994
1995
Asymmetry
Data - Fit
√s (GeV)

Results


ALEPH
DELPHI
L3
OPAL
LEP
0.0181 ± 0.0033
0.0189 ± 0.0048
0.0148 ± 0.0063
0.0069 ± 0.0051
0.0153 ± 0.0025
χ²/ DoF = 4.4/3
0
0.02
0.04
A^{0,e}_{fb}


ALEPH
DELPHI
L3
OPAL
LEP
0.0170 ± 0.0025
0.0160 ± 0.0025
0.0176 ± 0.0035
0.0156 ± 0.0025
0.0164 ± 0.0013
χ²/ DoF = 0.3/3
0.01
0.02
0.03
A^{0,μ}_{fb}


ALEPH
DELPHI
L3
OPAL
LEP
0.0166 ± 0.0028
0.0244 ± 0.0037
0.0233 ± 0.0049
0.0143 ± 0.0030
0.0183 ± 0.0017
χ²/ DoF = 5.9/3
0
0.01
0.02
0.03
0.04
A^{0,τ}_{fb}


ALEPH
DELPHI
L3
OPAL
LEP
0.0169 ± 0.0016
0.0187 ± 0.0019
0.0187 ± 0.0026
0.0141 ± 0.0017
0.0168 ± 0.0010
χ²/ DoF = 4.1/3
10³
10²
m_H [GeV]
0.01
0.015
0.02
0.025
0.03
A^{0,l}_{fb}

Lepton cross sections and asymmetries



Asymmetry at LEP2

Measurements continue at LEP2

- γ exchange dominates, $\sigma_{\ell\ell}$ reduced by factor ≈ 500, correspondingly less statistics.
- large fraction of 'radiative return' events
- analysis separated for inclusive ($\sqrt{s'/s} > 0.10$) and non-radiative ($\sqrt{s'/s} > 0.85$)





Asymmetry at LEP2



Results consistent with Standard Model expectations

Quark asymmetries

Hadronization obscures both flavour and charge of the primary quark.

Precise measurements possible for b and c

Two principal methods:

- particle tag (leptons, D-mesons) gives both quark flavour and charge.
- lifetime tag selects flavour,
  jet or hemisphere charge determines
  (statistically) primary quark charge.

$$Q_{jet} \equiv \frac{\sum_i Q_i |p_i^{||}|^\kappa}{\sum_i |p_i^{||}|^\kappa}$$

Lepton tags

Classical method to determine charge & and flavour of heavy quarks.

- Several sources must be distinguished



- Separation based on (p, p_T) of lepton w.r.t. jet axis



- Further improvement using additional info (lifetime, jet-shape) and neural net



- Uncertainties from
  - lepton spectra
  - leptonic branching ratios
  - mixing

  Best to determine $A^b_{\rm FB}$ and $A^c_{\rm FB}$ (+ mixing) in common fit.

	ALEPH	DELPHI	L3	OPAL
$A^b_{\rm FB}$	X	X	X	X
$A^c_{\rm FB}$		X		X
$\overline{\chi}$	X			X

$D^{(*)}$ reconstruction

$D^{(*)}$ mesons originate from two sources:

1. primary c quark + fragmentation quark
2. Deacy product of primary B meson

$\rightarrow$ both heavy flavour tag and charge id

D mesons reconstructed in various channels:

$$D^0 \rightarrow K^- \pi^+ \, , \; D^+ \rightarrow K^- \pi^+ \pi^+$$

$$\begin{aligned} D^{*+} \quad \rightarrow \quad & D^0 \pi^+ \\ & \;\; |\rightarrow K^- \pi^+ \\ & \;\; |\rightarrow K^- \pi^+ \pi^0 \\ & \;\; |\rightarrow K^- \pi^+ \pi^- \pi^+ \\ & \;\; |\rightarrow \ldots \end{aligned}$$

Slow pion characteristic signature for D^*:



D reconstruction used for $A_{\rm FB}^c$ by A D O.
Background from $B \to D$ decays controlled with D-momentum, lifetime, jet-shapes.

Jet charge

- Efficient & pure lifetime tags to select $b\bar{b}$ events.
- charge asymmetry

$$\langle Q_F - Q_B \rangle = \frac{N_F - N_B}{N_F + N_B} \langle Q_b - Q_{\bar{b}} \rangle = A_{\mathrm{FB}} \cdot \delta$$

- For a single flavour δ measured with data:

$$\delta \approx -4 \langle Q_F Q_B \rangle$$



- Correction to δ due to charge bias, background and correlations.
- Analogy to R_b double tag: Observable and main correction determined with data.
- Effects of QCD & $B^0\overline{B^0}$ mixing $\approx$ absorbed in δ
- Sophisticated techniques to extract maximal information:
  Fit as function of
  - b-purity
  - $\cos\theta$
  - κ ($Q_{jet} = \sum_i Q_i |p_i^{||}|^\kappa \,/\, \sum_i |p_i^{||}|^\kappa$)



A_{FB}^{b}

0.0625 ± 0.0192

0.1091 ± 0.0126

0.1021 ± 0.0094

0.0935 ± 0.0078

0.1041 ± 0.0063

0.1115 ± 0.0068

A^q_{FB} corrections

Measured A^q_{FB} $\rightarrow$ pure Z^0 $A^{0,q}_{\mathrm{FB}}$

- 'oblique' corrections due to ISR, γ, γZ and $\sqrt{s}$.
- Large QCD corrections ($\approx$ error), depending on details of analysis



$$A^q_{\mathrm{FB}} = (1 - B_{exp}\delta_{QCD})A^{0,q}_{\mathrm{FB}}$$

- δ_{QCD} analytically calculable ($\approx 5\,\%$)
- B_{exp} additional experimental bias, specific for experiment and method:

  leptons 0.5

  D recon. 0.2

  jet-charge < 0.1

Combination of heavy flavour results

Averaging heavy flavour results from different analyses & experiments complicated due to

- Correlations between e-weak observables (R_b, R_c)
- Correlations between physics input (mixing, production and decay rates)

LEP & SLD agreed on common definition of input parameters & errors.
Averaging done by fit to 88 input quantities from LEP & SLD
$\rightarrow$ 17 fit parameters.

$A_{\mathrm{FB}}^{b,c}$ energy dependence



Off-peak measurements consistent with SM expectation.

$A_{\mathrm{FB}}^{0,b}$ results



$A_{\mathrm{FB}}^{0,c}$ results



Tau polarization

τ only fermion where polarization precisely measureable.

- $(V-A)$ structure of τ charged current decay causes maximal parity violation.
- Decay directions in τ rest frame depend on polarization.

Most evident for $\tau \to \pi\nu$



$$\frac{dN}{d\cos\theta^*} \propto (1 + P_\tau \cos\theta^*)$$

$\cos\theta^*$ in τ rest frame related to E/E_{beam} in lab frame

$$\cos\theta^* \approx 2\frac{E_\pi}{E_{beam}} - 1$$

Other channels less direct:

- $\tau^- \to e^- \overline{\nu}_e \nu_\tau$ & $\tau^- \to \mu^- \overline{\nu}_\mu \nu_\tau$ are 3-body decays into 3 spin 1/2 particles, only 1 observed
  $\to$ signal strongly diluted

- Decays modes to spin-1 mesons (ρ, a_1) more complicated, two spin configurations of the meson. But subsequent decays ($\rho \to 2\pi$, $a_1 \to 3\pi$) give info on meson spin state.
  $\to$ part of τ polarization regained

	$\tau \to e\nu\nu$ $\tau \to \mu\nu\nu$	$\tau \to \pi\nu$ $\tau \to K\nu$	$\tau \to \rho\nu$	$\tau \to a_1\nu$
Br	0.35	0.12	0.25	0.09
Sens.	0.22	0.58	0.49	0.45
Weight	0.12	0.30	0.44	0.13

Polarization determined by fitting MC spectra for $\langle P_\tau \rangle = 1$ and $\langle P_\tau \rangle = -1$ to data.



Recent progress by new ALEPH analysis:
$\tau^+\tau^-$ direction determined for whole event:



τ direction given by intersection of cones (2 fold ambig.)

Improves sensitivity of a_1 and ρ channels
(0.48→0.58)



Angular distribution

$\langle P_\tau \rangle$ and $A_{\rm FB}^{\rm pol}$ determined by fitting observed $d\langle P_\tau \rangle / d\cos\theta$





Combined Results



Systematics mostly uncorrelated among experiments: Detector simulation, PID and calibration.

Correlated sources (a_1 decay, τ Br's, corrections) at the 1-2 % level.

Results by Channel

A_τ measurements by channel at LEP



π (23%)
ρ (28%)
a_1(3h) (88%)
e (82%)
μ (100%)

A_e measurements by channel at LEP



π (27%)
ρ (92%)
a_1(3h) (75%)
e (28%)
μ (21%)

Detector characteristics

Significant differences in precision already reflected in $\tau^- \to \nu_\tau \rho^-$ efficiency & purity.

	ALEPH	DELPHI	L3	OPAL
eff.	51 %	41 %	58 %	41 %
bkgr.	8.9 %	15 %	10.5 %	27 %

τ polarization poses specific challenges on ECAL performance. Most crucial π^0 id and separation of $\pi^0, \pi^\pm$ showers.

	ALEPH	DELPHI	L3	OPAL
Magnet	1.5 T	1.23 T	0.5 T	0.435 T
ECAL				
granularity	$3 \times 3 cm^2$	$2 \times 2 cm^2$	$2 \times 2 cm^2$	$10 \times 10 cm^2$
energy σ_E/E	$0.18/\sqrt{E}$	$0.32/\sqrt{E}$	$0.02/\sqrt{E}$	$0.06/\sqrt{E}$

Interpretation

Lepton universality:

Couplings $\hat{g}_v^l$ & $\hat{g}_a^l$ given by:

$\Gamma_l \propto (\hat{g}_v^l)^2 + (\hat{g}_a^l)^2$

$A_{\rm FB}^{0,l} \propto (\hat{g}_v^l/\hat{g}_a^l)^2$

$\langle P_\tau \rangle$ and $A_{\rm FB}^{\rm pol} \propto (\hat{g}_v^l/\hat{g}_a^l)$



$\sin^2\theta^{l}_{\mathrm{eff}}$



χ^2 quite ok now, was $12/6$ at EPS 1997

$(P(\chi^2) : 5\,\% \rightarrow 25\,\%)$

A_{LR} and $A_{\mathrm{FB}}^{0,b}$ disagree by about $2.5\ \sigma$.

$\rho^{\ell}_{\mathrm{eff}}$ vs $\sin^2\theta^{\,l}_{\mathrm{eff}}$

$$\Gamma_l \propto \rho = \frac{1}{1-\Delta\rho}$$



Presence of genuine electroweak corrections with >99 % C.L. (mainly due to ρ)

Pulls for SM fit



Overall χ^2/d.o.f. ok with $15/15$, no measurement more than $2\ \sigma$ away.
but $A^{0,b}_{\rm FB}$, A_{LR} & and $\mathcal{A}_b$ alone give $\chi^2 = 9.7$.

LEP and SLD asymmetries



Summary

- Lepton charge asymmetries essentially finalized. No change expected.
- Heavy quark asymmetries full statistics analyzed. Small improvements in some areas with refined analyses possible.
- Precision of LEP tau polarization substantially improved last year. Most analyses use full statistics. Some further gain in accuracy likely.
- Discrepancy in $\sin^2\theta^l_{\rm eff}$ measurements significantly reduced ($P(\chi^2) = 25\,\%$).

Standard Model still in perfect shape !